


Antipsychotic Utilization in a Pediatric Population

Megan J. Ehret, PharmD, MS, BCPP
 Professor; Co-Director of Mental Health Program
 University of Maryland, School of Pharmacy

1

My Perspective



4

Disclosures

- Dr. Ehret is a consultant for Saladex Biomedical. She serves as the Pharmacist Expert for SMI Adviser and is the Director for the State of Maryland Office of Pharmacy Services Peer Review Program.

2

The Approach

- Common case vignettes
- Tips
- Guidance
- Resources

5

Objectives

- Describe current practice guidelines regarding the use of antipsychotic medications in a pediatric population
- Outline adverse effects associated with the use of antipsychotic medication in a pediatric population
- Discuss when to initiate an antipsychotic medication in a pediatric population

3

Case 1


- 7 year old child with ADHD, PTSD, DMDD, MDD and unspecified anxiety. Provider has only seen child once.
- Current medications: Clonidine XR 1 mg daily
- Child referred to therapy
- Request for: Risperidone 0.5 mg daily
- Target symptoms: Aggression, assault, impulsivity, insomnia, irritability, mood instability

6

Poll Question

- Is Risperidone appropriate for this patient?
- Yes
- No
- I need more information

7



10

Antipsychotics in Pediatric Patients

Increasing in use since 1990

- Residential care > group homes > foster care > general population
- Publicly vs. privately insured

Most frequently prescribed

- Risperidone: 42.1%
- Aripiprazole: 28%
- Quetiapine: 19.2%
- Olanzapine: 4.4%

Most frequent target symptom: aggression

J Child Adolesc Psychopharmacol 2021;31:350-7
J Child Adolesc Psychopharmacol 2021;3:381-6
Front Psychiatry 2021;12:623681.ecollection 2021

8

What are we trying to do accomplish with ANTIPSYCHOTIC medication?

Achievable Therapeutic Objectives

- Enable the patient to control him/herself
- Relieve signs and symptoms which are troubling or disabling to the patient
- Improve the patient's ability for self-care

Antipsychotics don't control behavior

11

Prescribing Patterns

Age (Years)	Stimulant (%)	Antidepressant (%)	Antipsychotic (%)
3-5	0.5	0.2	0.2
6-12	4.6	1	0.8
13-18	3.7	2.8	1.2
19-24	1.6	4	0.8

- Highest prescribing among general psychiatry or child psychiatry for all age groups except stimulants
- Females prescribed antidepressants more than males as age increased
- Males aged 6-18 years prescribed antipsychotics and stimulants more than females

J Child Adolesc Psychopharmacol 2018;28:158-65

9

Psychiatric medication doesn't help


Developmental Issues:

- Learning disabilities (e.g. reading disability)
- Developmental delay (e.g. speech/language)
- Physical health (e.g. poorly controlled asthma)

Environmental Issues:

- School (e.g. frequent school changes)
- Family (e.g. parental illness or financial stressors)
- Community (e.g. bullying or violence exposure)

12



13

Assess Problem Behaviors

Timeline: episodic versus chronic

Settings: home, school, community

Impact: relationships, grades, safety

Context: stressors, changes

Emotion: irritability, anxiety

16

Tip #1:

Consider both the diagnosis and target symptoms.

14

Case 2

- 15 year old with depression and hallucinations
- Current medication: Quetiapine 75 mg daily, Clonidine 0.1 mg XR 2 daily, Escitalopram 20 mg daily, bupropion 100 SR daily, trazodone 50 mg daily
- Request for: Olanzapine 2.5 mg daily
- Target Symptom: Hallucinations

17

Pediatric Approved Antipsychotics

Antipsychotic	Irritability due to autism	Bipolar I	Schizophrenia
Aripiprazole (Abilify)*	X	X	X
Risperidone (Risperdal)	X	X	X
Olanzapine (Zyprexa)		X	X
Quetiapine (Seroquel)		X	X
Asenapine (Saphris)		X	
Paliperidone (Invega)			X
Lurasidone (Latuda)**		X	X


*Also has indication for treatment of Tourette's Disorder
 **Indication for bipolar depression

15

Poll Question

- Which recommendation is the best to provide for this patient?
 - Approve the olanzapine in combination with quetiapine
 - Decline the olanzapine, increase the quetiapine

18




19

Where else are Antipsychotics Recommended?

- Tourette’s Disorder
 - Second and third line options
- Autism
 - Risperidone and aripiprazole for irritability

22




20

Long-Acting Injectable Antipsychotic Medications- Pediatrics

- No prospective studies; majority case reports and case series
- Most reported LAI: Risperidal Consta, Invega Sustenna, Abilify Maintena
- Decrease in severity of symptoms and lower remission rates
- Possible decrease in AEs
- Barriers to administration
 - Providers knowledge
 - Clinic administration
 - Patient and guardian acceptance of LAI treatment
 - Insurance approval

J Child Adolesc Psychopharmacol 2022;32:312-27
Pediatric Drugs 2023;25:135-49

23



21

Case 3

- 14 year old with psychotic disorder NOS
- AIMS: 5
 - Lip Movement: 1
 - Upper Arm: 3
 - Lower Leg: 2
- CC: Tremors are not getting better
- Current Medications: Paliperidone 1.5 mg twice daily, benztropine 1 mg twice daily, lorazepam 1 mg three times daily
- Request: Paliperidone 1.5 mg AM, 3 mg PM
- Target Symptoms: Hyperactivity, impulsivity

24

Poll Question

- Should the dose of Paliperidone be increased?
 - Yes
 - No
 - I don't know what an AIMS score is

25

Adverse Effects

Evidence	<ul style="list-style-type: none"> •Short-term studies currently available •Need for long-term studies on brain development
Common Adverse Effects	<ul style="list-style-type: none"> •Somnolence/sedation •Gastrointestinal distress •Weight gain
Other Effects	<ul style="list-style-type: none"> •Metabolic Parameters- lipids, glucose, insulin resistance •Hormonal effects- prolactin •QTc Prolongation •NMS
Extrapyramidal Symptoms	<ul style="list-style-type: none"> •Decreased with SGA vs. FGA •Increased incidence in children and adolescents compared to adults

Psychiatr Serv 2016;67:339-41

28

Tip #3:
Safety first



26

Receptor Binding and Adverse Effects

	H ₁	H ₂	H ₃	M ₁	M ₃	α ₁	α _{2A}	α _{2B}	α _{2C}
Haloperidol		+				++	+	+	+
Aripiprazole	++					++	++	++	++
Asenapine	+++	+++				+++	+++	+++	+++
Lurasidone						++	++		+++
Olanzapine	+++	++	+	++	++	++	+	++	+++
Paliperidone	++	+				+++	+++	+++	+++
Quetiapine	+++			+	+	++	+	+	++
Risperidone	+++	+				+++	++	++	+++

Current Neuropharmacol 2018;16:1210-23

29

How do Antipsychotic Medications Differ?

- Side effect profile
- Efficacy
- Drug interactions
- Pharmacokinetics
- Dosage forms
- Regimen

27

Common Adverse Effect: Obesity

- Visible adverse effect: weight gain
- May not notice: Increased cholesterol, blood sugar
- Blood work and weight monitoring will improve early detection of problems
- Medication monitoring programs track side effect issues

30


Metabolic Side Effects

Antipsychotic	Weight Gain	Glucose Abnormalities	Hyperlipidemia
Aripiprazole	Low	Low	Low
Asenapine	Moderate	Moderate	Moderate
Brexipiprazole	Low	Low	Moderate
Cariprazine	Moderate	Moderate	Low
Clozapine	High	High	High
Haloperidol	Low	Low	Low
Iloperidone	Moderate	Moderate	Moderate
Lumateperone	Low	Low	Low
Lurasidone	Low	Moderate	Moderate
Olanzapine	High	High	High
Paliperidone	Moderate	Low	Moderate
Quetiapine	Moderate	Moderate	High
Risperidone	Moderate	Moderate	Low
Ziprasidone	Low	Low	Low

31

- ### Physical Morbidity and Mortality
- Sedentary lifestyle
 - Obesity
 - Cardiovascular diseases
 - Obstetric complications
 - Altered pain sensitivity
- *Lower rate of health care services utilization and medical treatment

34



32

Movement Disorders

- Acute Akathisia**
 - Complaints of restlessness
 - Excessive fidgety movements
 - Typically develop within a few weeks or starting medication or raising dose
- Acute Dystonia**
 - Severe muscle spasms of the eyes, head, neck, limbs, or trunk
 - Severe arching of the back or laryngospasm may occur
 - Symptoms typically emerge within a few days of starting medication or raising dose
- Parkinsonism**
 - Tremor
 - Muscular rigidity
 - Akinesia
 - Bradykinesia
 - Symptoms onset typically occurs within a few weeks to months of starting antipsychotic or raising dose
- Tardive Dyskinesia**
 - Involuntary, hyperkinetic, repetitive choreoathetoid movements
 - Most commonly observed in the orofacial region
 - Movements in extremities/trunk may also occur

CNS Spectrums 2015;20:4-14

35

Monitoring Metabolic Adverse Effects

	Baseline	4 Weeks	8 Weeks	12 Weeks	6 Months	Annually
Weight/ BMI	X	X	X	X	X	X
Fasting Plasma Glucose/ HgA1c	X			X	X	X
Lipids	X			X	X	X
Blood Pressure	X			X	X	X

J Psychopharmacol 2016;30:717-48
J Clin Psychopharmacol 2021;41:13-18

33

- ### Movement Disorders in Pediatrics
- Risk increased with duration of antipsychotic treatment:
 - 3% at 6-12 months
 - 10% at 1-2 years
 - 14% at more than 2 years
- Wonodi J, et al. Mov. Disord. 2007;22:1777-1782

36

Movement Disorders in Pediatrics

- 118 pediatric patients with psychiatric illness; 5-18 years treated for 6 or more months with antipsychotics
- >80% of antipsychotic prescriptions were for youth with no psychotic symptoms (mood disorders; ADHD)
- 9% showed TD

Wonodi I, et al. Mov. Disord. 2007;22:1777-1782

37

Safety of Long-Acting Injectable Antipsychotic Medications

40

Monitoring for Movement Disorders

Disorder	Monitoring Recommendations
Acute Dystonic Reaction	Simpson-Angus Extrapyramidal Symptom Scale Extrapyramidal Symptom Rating Scale Observation and patient report at each visit
Parkinsonism	Simpson-Angus Extrapyramidal Symptom Scale Extrapyramidal Symptom Rating Scale Observation and patient report at each visit
Akathisia	Barnes Akathisia Rating Scale Ask patient about this side effect and suicidal thoughts/plans at each visit
Tardive Dyskinesia	Abnormal Involuntary Movement Scale (AIMS) or Dyskinesia Identification System-Condensed User Scale (DISCUS) at baseline and at least every 6 months/12 months

38

Case 4

- 16 year old with Bipolar Disorder and intellectual disability
- Current Medication: Olanzapine 10 mg daily
- Request: Zyprexa Relprevv LAI
 - Will have PCP administer injection and then patient will come to psychiatry provider office for the 3 hour monitoring

41

AIMS Rating Scale

Patient Name:	Date:	None	Mild/moderate	Mild	Moderate	Severe
Facial and Oral Movements						
1. Muscles of facial expression (e.g., movements of forehead, eyebrows, periorbital area, cheeks). Include frowning, blinking, smiling, grimacing.						
0	1	2	3	4		
2. Lips and perioral area (e.g., puckering, protruding, smacking)						
0	1	2	3	4		
3. Jaw (e.g., biting, clenching, chewing, mouth opening, lateral movement)						
0	1	2	3	4		
4. Tongue (Rate only increases in movement both in and out of mouth; NOT inability to sustain movement)						
0	1	2	3	4		
Extremity Movements						
5. Upper (arms, wrists, hands, fingers) include choreic movements (i.e., rapid, objectively purposeless, irregular, spontaneous) athetoid movements (i.e., slow, irregular, complex, serpentine). (DO NOT include tremor (i.e., repetitive, regular, rhythmic).)						
0	1	2	3	4		
6. Lower (legs, knees, ankles, feet) (e.g., lateral knee movement, foot tapping, heel dragging, foot spurning, inversion and eversion of foot)						
0	1	2	3	4		
Trunk Movements						
7. Neck, shoulders, hips (e.g., rocking, twisting, spurning, pelvic gyrations)						
0	1	2	3	4		
Global Judgments						
8. Severity of abnormal movements						
0	1	2	3	4		
9. Interference due to abnormal movements						
0	1	2	3	4		
10. Patient's awareness of abnormal movements (rate only patient's report)						
0	1	2	3	4		
0 = not aware; 1 = aware, no distress; 2 = aware, mild distress; 3 = aware, moderate distress; 4 = aware, severe distress						
Dental Status						
11. Current problems with teeth and/or dentures?						
No	Yes					

39

Poll Question

- Does this plan meet the requirements of the REMS for Zyprexa Relprevv?
 - Yes
 - No
 - I have no idea

42

Tip #4:

Have a consultation plan for outliers



43

Injection Sites/Volume of Injection

Medication	Site	Volume of Injection
Risperdal Consta	Deltoid or Gluteal IM	2 mL
Perseris	Abdomen or back of upper arm subcutaneously	0.6-0.8 mL
Rykindo	Gluteal IM	2 mL
Uzedy	Abdomen or back of upper arm subcutaneously	0.14-0.7 mL
Zyprexa Relprevv	Gluteal IM	1-2.7 mL

46

Injection Sites/Volume of Injection

Medication	Site	Volume of Injection
Fluphenazine	Typically IM, with deltoid or gluteal injections possible	0.5-2 mL
Haloperidol	Deltoid or gluteal IM	3 mL max

44

Safety Concerns

- Needle Size
 - Consideration of body mass
 - Adult studies versus pediatric patients
- Preparation
 - Training to avoid errors
- Oral overlap
- Long-term data not available in pediatric patients

47

Injection Sites/Volume of Injection

Medication	Site	Volume of Injection
Abilify Maintena	Deltoid or Gluteal IM	0.8-2 mL
Abilify Asimtufii	Gluteal IM	3.2-2.4 mL
Aristada	441 mg: Deltoid or Gluteal IM Higher doses: Gluteal IM	1.6-3.9 mL
Aristada Initio	Deltoid or Gluteal IM	2.4 mL
Invega Sustenna	Loading dose: Deltoid IM Maintenance Dose: Deltoid or Gluteal IM	0.25-1.5 mL
Invega Trinza	Deltoid or Gluteal IM	0.875-2.625 mL
Invega Hafyera	Gluteal IM	3.5-5 mL

45

Zyprexa Relprevv

- REMS: Potential for development of Post-Injection Delirium/Sedation Syndrome (PDSS)
 - Signs and symptoms consistent with olanzapine overdose, in particular sedation and/or delirium
 - Must be administered in registered healthcare facility with ready access to emergency response services
 - Continuously observe patient for at least 3 hours
 - After 3 hours, confirm patient is alert, oriented, and absent of any signs/symptoms of PDSS
 - Must have ride to next location

48

Resources

- Movement Disorder Training
 - The AIMS Assessment and Tardive Dyskinesia
- Psychotropic Long-Acting Injectable (LAI) Training Program
 - Psychotropic LAI Training Program

49

Questions

52

SMI ADVISER APP



50

50

Conclusions

- Verify indications and target symptoms
- Most common adverse effects: hypotension, increased appetite, weight gain, and sedation
- Changes in glucose and lipids
- Treatment plans should be based on treatment guidelines

51